



US005171671A

United States Patent [19]

Evans et al.

[11] **Patent Number:** 5,171,671[45] **Date of Patent:** Dec. 15, 1992[54] **RETINOIC ACID RECEPTOR COMPOSITION**

[75] **Inventors:** Ronald M. Evans, La Jolla; Estelita S. Ong; Prudimar S. Segui, both of San Diego; Catherine C. Thompson, La Jolla; Kazuhiko Uemsono, San Diego, all of Calif.; Vincent Giguere, Etobicoke, Canada

[73] **Assignee:** The Salk Institute for Biological Studies, San Diego, Calif.

[21] **Appl. No.:** 546,256

[22] **Filed:** Aug. 6, 1990

Related U.S. Application Data

[60] Division of Ser. No. 276,536, Nov. 30, 1988, Pat. No. 4,981,784, which is a continuation-in-part of Ser. No. 128,331, Dec. 2, 1987, abandoned.

[51] **Int. Cl.⁵** C12N 15/12; C12N 15/62; C12N 15/63; C07K 13/00

[52] **U.S. Cl.** 435/69.1; 435/69.7; 435/257.3; 435/320.1; 530/350; 536/27

[58] **Field of Search** 536/27; 435/69.1, 69.7, 435/252.3, 320.1; 530/350

[56] **References Cited****PUBLICATIONS**

Nature 332:850-853, Apr. 28, 1988, Brard et al., Identification of a second human retinoic acid receptor.
 Science 240:889-895, May 13, 1988, Evans, The Steroid and Thyroid Hormone Receptor Superfamily.
 Nature 330:624-629, Dec. 17, 1987, Giguere et al., Identification of a receptor for the morphogen retinoic acid.
 Nature 330:444-450, Dec. 17, 1987, Petkovich et al., A human retinoic acid receptor which belongs to the family of nuclear receptors.

Nature 330:420-421, Dec. 3, 1987, Robertson, Towards a biochemistry of morphogenesis.

Pharm. Rev. 36:935-1005, 1984, Chytil, Retinoic Acid: Biochemistry, Pharmacology, Toxicology, and Therapeutic Use.

PNAS, 84:5645-5649, Aug. 1987, Shubeita et al., Molecular cloning and analysis of functional cDNA and genomic clones encoding bovine cellular retinoic acid . . .

Primary Examiner—David L. Lacey

Assistant Examiner—John D. Ulm

Attorney, Agent, or Firm—McCubbrey, Bartels, Meyer & Ward

[57]

ABSTRACT

A novel retinoic acid receptor is disclosed. The novel receptor is encoded for by cDNA carried on plasmid pHRAR1, which has been deposited with the American Type Culture Collection for patent purposes. Chimeric receptor proteins are also disclosed. The chimera are constructed by exchanging functional domains between the glucocorticoid, the mineralocorticoid, the estrogen-related, the thyroid and the retinoic acid receptors. In addition, a novel method for identifying functional ligands for receptor proteins is disclosed. The method, which takes advantage of the modular structure of the hormone receptors and the idea that the functional domains may be interchangeable, replaces the DNA-binding domain of a putative novel receptor with the DNA-binding domain of a known receptor such as the glucocorticoid receptor. The resulting chimeric construction, when expressed in cells, produces a hybrid receptor whose activation of a ligand (e.g., glucocorticoid) inducible promoter is dependent on the presence of the new ligand. The novel method is illustrated in part by showing that the ligand for the new receptor protein is the retinoid, retinoic acid.

14 Claims, 12 Drawing Sheets

J1046 U.S. PTO
 09/773042
 01/31/01



Patent and Trademark Office